

Thermal Equilibrium

Basics

- ▶ **Thermal equilibrium** is a central concept in [thermodynamics](#). It describes the unique state of an ensemble of particles (i.e. the atoms of a crystal) that the system assumes by itself sooner or later (and later can mean really, really late) for a given set of intrinsic parameters (e.g., temperature, pressure, [chemical potential](#)) and extrinsic parameters (e.g., volume, entropy, number of particles).
 - The state of the system is unambiguously described by a state function which is called a **thermodynamic potential** and there are several thermodynamic potentials that can be used for a system description.
 - Whereas in principle *any* thermodynamic potential can be used for *any* situation (because they are related by a so-called **Legendre transformation**); it is useful to use specific thermodynamic potentials for specific systems.
- ▶ Depending on the kind of "contact" between the system under consideration and the environment (e.g. totally isolated, energy flow permitted, particle flow permitted, and so on), typical situations are:
 - ▶ **Constant volume V , temperature T , and number of particles N .**
 - The proper thermodynamic potential is the **free energy $F(V, T, N)$** (sometimes called **Helmholtz energy**).
 - ▶ **Constant pressure p , constant temperature T , and constant particle (= atom) number N**
 - This is the situation typical for a crystal. The appropriate thermodynamic potential is the **free enthalpy $G(p, T, N)$** (sometimes called **Gibbs energy**).
 - ▶ The **free enthalpy** (defined as **$G = H - TS$**) with **$H =$ enthalpy** of the system and **$S =$ entropy** is thus the most important thermodynamic potential when considering defects.
 - Thermal equilibrium for this case then simply means a state with an (absolute) minimum of the free enthalpy of the crystal.