

Boltzmann Distribution

Classical Particles

Science

Let's give a quick look to one of the deepest and most useful insights in thermodynamics. In order to start simple, I will first go back to the good old times when quantum theory and so on had not yet been discovered, and particles like atoms, molecules and so on were at best "classical" particles or mass points - just very small.

Classical physics deals with those mass-points by using *classical* mechanics, going back all the way to **Newton**.

As long as your "particles" - a cannon ball, a stone, the earth and the sun, a hydrogen atom - could be described as a "points" positioned at the center of gravity, all is well. You could precisely account - in principle - for the motion of the sun around the earth, or the movement of balls on the pool table, even when the particles occasionally hit each other. The only problem was that math or computers were not evolved enough to really let you do the calculations. There is a big difference between writing down a bunch of equations and actually solving them. But even if *you* couldn't do the required math, there wasn't a sliver of doubt that the solution, if obtained by somebody else, would tell you precisely what will happen.

Now make your pool balls much smaller, increase their number to 10^{24} , and allow them to move in three directions instead of just two. There was *no way* then, and there is still *no way* now, to calculate anything by following the movement of *each and every* ball.

Of course, a *pool* game like this does not exist, so why worry? Because *systems* like this do exist under different names. They are called, for example:

- the air in the room you're breathing at this very moment.
- crystals, like iron or silicon.
- your sword.
- you.

All those systems consist of a large bunch of atoms = particles that can move, more or less restrictedly, in three directions. Even the atoms in crystals or in your body can move a bit: they vibrate in three directions.

Dealing with systems like that (and you realize now that this happens a hell more often than dealing with systems that can be described by just a few mass points) necessitates to look at *statistical parameters* of the particle properties and their doings; things like *averages* or standard deviations.

The branch of physics dealing with that is called **statistical thermodynamics**, and that part of physics, together with quantum theory, turns chemistry into a branch of physics.

What kind of statistical properties am I talking about? At least two of those properties are well known to all of us, even to those who couldn't even spell "statistics":

1. The (unordered) *average energy* of a particle in a system of many. The common word for that is **temperature**.
2. The *average momentum* of a particle in a system of many. The common word for that is **pressure**.

So far so easy. If we now go one step deeper, we don't just look for averages but for the **distributions** of the parameters in question.

Averages are fine but do they not tell you all that much about your system. You could get an energy average of "5" for a system with 100 particles if, for example:

- 99 particles have a energy of "0" and one has the value 500.
- All have energy = 5.
- 50 have energy = 6, and the other 50 energy = 4.
- You get the point, pick your own example.

There are plenty of other ways or *distributions* of the energy values that give the same average. Some distributions are more orderly than others; the one with "all have energy = 5" certainly has a high degree of order. That's obviously where the [second law](#) and [entropy](#) comes in. It demands categorically that out of the large number of possible distributions that give the same average, only those that comply with the second law will really occur in nature.

That tells us what to do. Calculate energy and entropy for all possible distributions, derive the free enthalpy from that and see which distributions gives you the lowest value. That will be the one Mother Nature will implement.

The only problem is that going through the whole program of calculating entropy and free energy, finding the minimum, and so on, isn't all that easy, even for the most simple case of [vacancies in a crystal](#).

Enter **Ludwig Boltzmann**. He (with some help from others) simplified everything to a just amazing degree.

"You are looking at a (classical) system, any system, where the available energies are known", he says, "and you want to know how many particles have the energy E in a system with N_0 particles *that is in equilibrium at the temperature T* ? Well, here is the answer:"

$$\frac{N}{N_0} = \exp - \frac{E}{kT}$$

Boltzmann distribution or Boltzmann statistics (approximation)

- This is good for all cases where the energy E considered is substantially larger (let's say at least 0.1 eV) than E_0 , the smallest possible energy always set at $E_0 = 0 \text{ eV}$.

There are some minor details to consider if one goes for full generality but that is of no interest here.

- The Boltzmann distribution is a rather simple but *extremely* powerful equation. The exponential term is called "**Boltzmann factor**".

Application of Boltzmann Distribution to Diffusion

- Now let's see what the Boltzmann distribution can do for us. First, we notice that with its help we can write down the vacancy concentration, that took us a lot of effort to derive, just so. All we need to do is to reformulate the problem.

- Whenever an atom in a crystal "has" the energy E_V = formation energy of a vacancy, it will "somehow" escape and a vacancy is formed. The "somehow" doesn't need to worry us. In equilibrium, whatever was needed to be done to achieve it (here: get the atom out), has been done in one way or other, we wouldn't have equilibrium otherwise.

So how many atoms = vacancies do we have at the energy of E_V , the formation energy of a vacancy? Use the Boltzmann distribution from above and you get

$$N_V = N_0 \cdot \exp - \frac{E_V}{kT}$$


Vacancy concentration from Boltzmann distribution

- That's exactly what we got stomping through a lot of tricky math! Isn't that something?
- Thinking a little bit about all that, we realize that the Boltzmann *factor* simply describes the **probability** for some particles that an event needing the energy E will occur.

That is the best way to interpret the Boltzmann factor, and that will be very helpful right below.

- Now let's go for something else. We wonder how often a vacancy makes a jump, or, more precisely, how often one of the neighboring atoms jumps into the vacancy. Here is the link to a picture and the other stuff concerning this issue, which is of overbearing importance to sword making (and about anything else).

- In a somewhat more abstract way the situation is as follows:



- The blue atom (as all the other atoms) vibrates about its average position with a [typical vibration frequency](#) of $\nu \approx 10^{13} \text{ Hz}$ as schematically indicated.

The blue line indicates the potential energy of the atom. Only if the atom has enough energy (= large vibration amplitude) to overcome the energy barrier between the two positions, can it jump into the vacancy, ending up in a neighboring position. In the figure the height of the energy barrier is $E_2 - E_1 = E_M$; a quantity that is typical for a given crystal and called **migration energy** E_M .

So how often does an atom next to a vacancy jump into the vacancy? Easy. How often do you jump across a hurdle with a certain height? It is simply the number of attempts per time unit *times* the probability that a try was successful. For the atom the number of tries is just its vibration frequency ν (that is how often it runs up the barrier per second). Multiply with the probability that it has the required energy E_M (= Boltzmann factor for that energy). In math shorthand we have

$$r = \text{jump rate} = \nu \cdot \exp\left(-\frac{E_M}{kT}\right)$$

- How many atoms *in total* will jump per second? Well, only *one* of the atoms bordering a vacancy gets a chance for a jump. So the number or rate R of *all* atoms jumping per second it just the number of vacancies or $N_V = N_0 \nu \cdot \exp\left(-\frac{E_M}{kT}\right)$ *times* the rate of jumps for any of those atoms, or

$$\frac{R}{N_0} = \nu \cdot \exp\left(-\frac{E_M}{kT}\right) \cdot \exp\left(-\frac{E_V}{kT}\right) = \nu \cdot \exp\left(-\frac{(E_M + E_V)}{kT}\right)$$

- $(R/N_0) \cdot 100$ would be the percentage of the atoms in the crystal that jump per second. If we want to be absolutely precise, we have to make some extremely minor adjustments to this equation, but the essence of all this is:

We have diffusion in crystals covered!

I could go on and on about what you can do with the Boltzmann distribution. It certainly is one of the most important relations in Materials Science and there are plenty of equations that contain a *Boltzmann factor*. I won't. I rather give you a very brief and supercilious glance on what happens for *real* particles. The Boltzmann distribution, after all, is only valid for *classical* particles that do not really exist but are only *approximations* to the real thing.

Boltzmann Distribution and "Real" Particles

Once more: There is no such thing as a **classical particle** that follows Boltzmann statistics. Sorry. Classical particles, it turns out, are just **approximations** to *real* particles.

- As you (should) know, there are many different kinds of [real particles](#) out there. As you also know, there are many flavors of "ice cream" out there - but just two basic kinds: "cream" and "sorbet". Same thing for wine; you have red or white. People, including extremely strange guys like (*insert your choice*), come always as male or female.

While there are many kinds of particles, including very strange stuff like "naked bottom quarks" and simple things like protons, neutrons, electrons, atoms, photons, positrons, muons, ..., they also come in just *two* basic kinds:

- **Fermions.** Fermions must never ever have the same "state".
Examples: Electrons, protons, ..
- **Bosons.** Bosons are allowed to have the same state.
Examples: Photons, some mesons, combinations of fermions; e.g. some atoms, ..

- So what is a (quantum mechanical) "state"? It is simply *one* of the solutions of the [Schrödinger equation](#), the fundamental equation that replaces "Newton's laws" in quantum mechanics, for a given problem. The problem could be the bunch of electrons careening around in a metal, or a lot of atoms vibrating in a crystal, or anything else you care to formulate.
For general problems like that there are always *infinitely* many solutions - but that is nothing new. If you calculate for a classical particles what kind of velocities or energies they *could* have, the solution of the problem says: "infinitely many; *any* you like are possible" (there are no restrictions of velocities with only "Newton"!). The *real* problem in somewhat more advanced science / math is always to find the *particular* solutions, out of many, that apply to a particular sub group of the problem.

In quantum mechanics, we have infinitely many solutions to the question above as in classical mechanics, *but not all velocities and energies are allowed*. Allowed energies, to give a (schematic) example, might be $E = 1, 2, 3, ..$ but never $E = 1.2$ or $E = 3.765$.


- To stay simple, let's assume for the moment that every state = special solution of the Schrödinger equation has exactly *one* energy associated with it, and that every state has a *different* energy.
That allows us to look in a simple way on *quantum* distribution functions. We do that separately for Bosons and Fermions.

1. Distribution function for Bosons or **Bose-Einstein statistics**.

- All we need to know is:
 - The exact equation is somewhat different from the Boltzmann statistic [as given above](#).
 - In most practical cases the mathematically simpler Boltzmann statistics is a good approximation to the Bose-Einstein statistics.
This is as it should be. In both cases an arbitrary number of particles could have the same energy. In particular, all particles *could* have the lowest energy, and at very low temperatures all particles *will* have that lowest energy.
- I hear you. "What exactly is then the difference between a classical particle, that you claim does not exist, and a Boson? It looks as if they do pretty much the same thing".
Well, yes - but there *is* a difference. *Classical* particles we can always *distinguish* in principle. We can, so to speak, attach a number, paint them in different colors, or give them a name to which they respond. No matter how small they are, there is always something smaller (your classical particles divided into 1.000 smaller ones, for example), that you can use to make some kind of label.
Real particles, no matter if they are bosons or fermions, we *cannot* distinguish. There is no way to figure out if the particular electron that just produced some current in your solar cell is the same one that did it 5 minutes ago. And it is not that we just lack the means to do so. It is an intrinsic or inalienable property of real particles that they are indistinguishable. The universe subscribes to communism: all particles (of one kind) are exactly equal and indistinguishable. No proton will ever be richer, taller, healthier, more beautiful, smarter, or ([insert your choice of properties](#)) than all the other ones.
- If you think about it a for a few years, you start to realize that this is really, really weird. But that is the way the universe works and that's why there is a certain (if often small) difference between classical particles and Bosons.

2. Distribution function for Fermions or **Fermi-Dirac statistics**.

- Now we have a huge difference to classical particles. It's best to look at a basic figure to see that.



- Shown is the *number* of particles at a certain energy for the Boltzmann case, and the *probability* that a particle is found at a certain energy for the Fermi case.
We need two different scales because on just one scale one of the two curves would disappear in an axis and become "invisible".
- In the *Boltzmann case* we have a hell of a lot of a particles with energies at or close to the minimum energy of the system (always at $E_0 = 0 \text{ eV}$). The number "10²⁰" on the scale has no specific meaning and only illustrates "a hell of a lot". If the temperature is decreased, the curve gets steeper; at $T = 0 \text{ K}$, it is identical with the axis

- In the *Fermi case* we have a probability of **1** for lower energies that a particle "has" that energy or, as we say, that the place at some allowed energy is occupied by one of the fermions of the system. This goes up to very high energies since all particles need to occupy some place, and so far only one particle is allowed per energy level. At **T = 0 K** the curve would be a rectangular box. The particles of the system occupy as many energy levels as are needed to accommodate them all. Occupation ends only around some special energy called **Fermi energy E_F** , where at **T = 0 K** the last particle finds its place. Places above the Fermi energy then are empty. At *finite* temperatures, the Fermi distribution "softens" around the Fermi energy as shown. A few particles have higher energies than the Fermi energy, and some places at lower energies are unoccupied. That produces some disorder or *entropy*, needed for nirvana or thermal equilibrium since the *second law* still applies.
 - For most applications, only the "high-energy tail" of the Fermi distribution is important. This part can be approximated very well with the (much simpler) Boltzmann distribution as shown. In other words: High energy fermions / electron behave pretty much like *classical* particles.
- And now we know (sort of) under what kinds of circumstances real particles might be approximated by classical particles. This is great because the math for classical particles is so much easier than that for real particles. But that is the *only* reason why we use the Boltzmann distribution and so on.
- For the finish, let's look at some of the most important *equations* for electrons; our most important fermions:

$$n(E; E_F, T) = D(E) \cdot f(E; E_F, T) \cdot \Delta E$$

$$f(E; E_F, T) = \frac{1}{\exp\left(\frac{E - E_F}{kT}\right) + 1}$$

Fundamental Equations for Electronics

$f(E; E_F, T)$ = Fermi distribution.
 $n(E; E_F, T)$ = concentration of electrons at energy E
 $D(E)$ = "density of states".

- Those are two of the few fundamental equations for all of solid state electronics. They are at the root of almost everything - from the simple conductivity of a copper wire to the transistors and so on in microchips. And they are far simpler than there looks (a bit like Wagner's music that is far better than it sounds according to Mark Twain). Let's see:
- $f(E; E_F, T)$, the Fermi-Dirac distribution, is a function of the energy E , with Fermi energy E_F and temperature T as *parameters*. The Fermi *distribution* is a universal function, it is valid for all fermions everywhere and everywhere. The Fermi *energy*, however, is a kind of material constant and thus describes some property of a given material. The "shape" or graph of the Fermi distribution is simple, it looks as shown above.
 - Now we must overcome our "one electron - one energy" restriction from above. In all *real* systems, for example the system of the electrons in a piece of silicon (Si), there are always several states that happen to have the same energy. Since electrons only must have different states, we can have as many electrons with same energy as there are states to that energy. The number of states with the same energy we call "**density of states**" $D(E)$. The number or density (number per volume) of electrons within some small energy interval ΔE is then simply given by multiplying the number of places there (= $D(E) \cdot \Delta E$) with the probability $f(E; E_F, T)$ that a place is occupied. That is what the first equation specifies. It also gives us the density of electrons in semiconductor that we can "work with" and thus the most important parameter for all semiconductor devices.

Looking at all of this in a general way, we see how technology emerges out of science:

- The Fermi distribution $f(\mathbf{E}; \mathbf{E}_F, T)$ is a universal (= valid in the whole universe) and rather simple function. It expresses a basic property of the universe and thus rather deep *science*.
- The density of states $D(\mathbf{E})$ is material specific and typical for the *perfect* crystal. It is always a rather complicated function, hard to calculate but not too difficult to measure. It expresses basic electronic properties of an *ideal crystal* and cannot be changed.
- The Fermi energy \mathbf{E}_F , while linked to the density of states, is sensitive to crystal *defects*. It therefore expresses properties of a *real crystal* and can be changed.
- The temperature T is "the *agent of change*". It changes the Fermi distribution and the Fermi energy to some extent.

Where does that leave you when you want to make an electronic device, implying that you must have the "right" electron density n ?

Well, provided you know what your device is supposed to do, you go through these steps:

- Define the density of states you need. In other words. Pick the best suited semiconductor.
- Manipulate the Fermi energy in all or in parts of your material (= semiconductor) by introducing the right defects at the right places. Most people call that procedure "doping".
- Use temperature to achieve proper distributions of the doping defects (= substitutional impurities).

Easy. Well, not quite. There are a few other things you need for making devices, like insulators and conductors, besides your semiconductor. Unfortunately, the "defect engineering" required to adjust the Fermi energies, is not all that simple. It can be done, however, as the computer you are using right now amply demonstrates.