

# Required Reading


## 1.3.5 Averaging Vectors

Let's look a bit closer at the averages one can take by considering a (localized) swarm of summer flies "circling" around like crazy, so that the ensemble looks like a small cloud of smoke.

- "Localized" means that the swarm maintains a defined form (on average), so all flies are always inside this defined volume in space.
- In the case of the charge carriers in a piece of metal, it simply implies that the carriers stay inside the piece of metal.

First we notice again that while the *individual* fly moves around quite fast, its average *vector velocity*  $\langle \underline{v}_i \rangle_t$ , averaged over time  $t$ , must be zero as long as the swarm as an ensemble doesn't move.

- In other words, the flies, on average, move just as often to the left as to the right etc. The net current flowing through some surface produced by *all flies at any given instance*, or by *one individual fly after sufficient time* is obviously zero for *any* reference surface you care to chose. This is illustrated schematically below.



- Shown are 13 velocity vectors of an individual fly; the chain of vectors closes so  $\langle \underline{v}_i \rangle_t = \mathbf{0}$ .

The average of the *magnitude* of the velocity of an *individual* fly,  $\langle |\underline{v}_i| \rangle_t = \langle \underline{v}_i \rangle_t$ , however, is obviously *not* zero - the fly, after all, is buzzing around at high (average) speed. *Note the details in the equation above: Only the underlining of v is different!*

- If we define  $\langle \underline{v}_i \rangle_t$  as follows, we have a simple way of obtaining the average of the magnitude (we take only the positive root, of course) .

$$\langle \underline{v}_i \rangle_t = + \langle \underline{v}_i^2 \rangle_t^{1/2}$$

- $\underline{v}^2$  is a scalar, and the (positive) square root of  $\underline{v}^2$  gives always the (positive) magnitude of  $\underline{v}$ ; i.e.  $|\underline{v}|$

- This is an elegant and workable definition, but beware:

$\langle \underline{v}^2 \rangle^{1/2}$  is *not* the same as  $(\langle \underline{v}^2 \rangle)^{1/2}$ !

Lets try it with a few arbitrary numbers →

$ \underline{v}  =$	3	4	6
$\langle \underline{v}^2 \rangle^{1/2} =$	$(3 + 4 + 6)/3 = 13/3 = 4,333\dots$		
$(\langle \underline{v}^2 \rangle)^{1/2} =$	$[(9 + 16 + 36)/3]^{1/2} = 20,33^{1/2} = 4,51$		

If we have  $\langle \underline{v} \rangle_t = \langle \underline{v}^2 \rangle_t^{1/2}$ , we may also calculate the average (over time) of the velocity *components* in *x*, *y*, and *z*-direction,  $\langle \underline{v}_x \rangle_t$ ,  $\langle \underline{v}_y \rangle_t$ ,  $\langle \underline{v}_z \rangle_t$ , of an individual fly *for a truly random movement*. (*We drop the index "i" now to make life easier*).

- Again, the vector averages  $\langle \underline{v}_x \rangle$  and so on of the *vector* components must be =  $\mathbf{0}$  because in a truly random movement the components in + *x* and - *x* direction and so on must cancel on average.
- Since the magnitude  $|\underline{A}|$  of a vector **A** is given by the square root of the scalar product of the vector with itself . We have

$$\underline{A} \cdot \underline{A} = A_x \cdot A_x + A_y \cdot A_y + A_z \cdot A_z = A^2$$

$$A = |\underline{A}| = (A^2)^{1/2}$$

Since

$$\langle v^2 \rangle_t = \langle v_x^2 \rangle_t + \langle v_y^2 \rangle_t + \langle v_z^2 \rangle_t ,$$

and since in a *truly random movement* we have

$$\langle v_x \rangle_t = \langle v_y \rangle_t = \langle v_z \rangle_t ,$$

we end up with

$$\langle v^2 \rangle_t = 3 \langle v_x^2 \rangle_t$$

$$\langle v_x^2 \rangle_t = 1/3 \langle v^2 \rangle_t .$$

From this we finally get

$$\langle v_x \rangle_t = \langle (v_x^2)^{1/2} \rangle_t = (1/3)^{1/2} \cdot \langle (v^2)^{1/2} \rangle_t = \frac{\langle v \rangle_t}{3^{1/2}}$$

In real life, however, the fly swarm "cloud" often moves slowly around - it has a finite **drift velocity**  $\underline{v}_D$ .

$$\underline{v}_D = \langle \underline{v}_i \rangle_t$$

- In consequence,  $\langle \underline{v}_i \rangle_t$  is not zero, and  $\langle \underline{v}_i, +x \rangle_t$  (= average velocity component in  $+x$  direction) in general is different from  $\langle \underline{v}_i, -x \rangle_t$ .
- Note that the drift velocity by definition is an average over vectors; we do not use the  $\langle \rangle$  brackets to signify that anymore. Note also, that the drift velocity of the *fly swarm* and the drift velocity of an *individual fly* must be identical if the swarm is to stay together.
- Without prove, it is evident that  $\underline{v}_D, i, x = \langle \underline{v}_i, +x \rangle_t - \langle \underline{v}_i, -x \rangle_t$  and so on. In words: The magnitude of the component of the average drift velocity of fly number  $i$  in  $x$ -direction is given by the difference of the average velocity components in  $+x$  and  $-x$  direction.

This induces us to look now at the *ensemble*, the swarm of flies. What can we learn about the averages taken for the *ensemble* from the known averages of *individual* flies?

- As long as every fly does - on average - the same thing, the *vector* average over time of the ensemble is identical to that of an individual fly - if we sum up a few thousand vectors for *one* fly, or a few million for *lots* of flies does not make any difference. However, we also may obtain this average in a different way:
- We do not average *one fly in time* obtaining  $\langle \underline{v}_i \rangle_t$ , but at any given time *all flies in space*.
- This means, we just add up the velocity vectors of all flies at some moment in time and obtain  $\langle \underline{v}_e \rangle_t$ , the **ensemble average**. It is evident (but not easy to prove for general cases) that

$$\langle \underline{v}_i \rangle_t = \langle \underline{v}_e \rangle_t$$

- i.e. *time average = ensemble average*. The new subscripts "e" and "r" denote ensemble and space, respectively. This is a simple version of a very far reaching concept in stochastic physics known under the catch word "**ergodic hypothesis**".

This means that in "normal" cases, it doesn't matter how averages are taken. This is the reason why text books are often a bit unspecific at this point: It is intuitively clear what a drift velocity is and we don't have to worry about how it is obtained. It also allows us to drop all indices from now on whenever they are not really needed.

- In our fly swarm example, the drift velocity  $\langle \underline{v}_D \rangle = \langle \underline{v}_i \rangle$  is usually much smaller than the average  $\langle v_i \rangle$  of the velocity magnitudes of an individual fly.

- The magnitude of  $\langle v_D \rangle$  is the difference of two large numbers - the average velocity of the *individual* flies in the drift direction *minus* the average velocity of the *individual* flies in the direction opposite to the drift direction.
- This induces an *asymmetry*: From a knowledge of the drift velocity *only*, *no inference whatsoever* can be made with regard to  $\langle v_i, +x \rangle$ ,  $\langle v_i, -x \rangle$  or  $\langle v_i \rangle$  whereas knowledge of  $\langle v_i, +x \rangle$  and  $\langle v_i, -x \rangle$  tells us all there is to know in *x*-direction

▀ This teaches us a few things:

- 1. Don't confuse  $\langle v \rangle$  with  $\langle v \rangle$ . The first quantity - for our flies - is zero or small, whereas the second quantity is large; they are totally different "animals".
- 2. This means in other words: Don't confuse the property of the *ensemble* - the drift velocity  $v_D$  of the ensemble or swarm - with the properties of the *individuals* making up the ensemble.