

## Solution to Exercise 1.3-2

### Derive numbers for $v_0$ , $v_D$ , $\tau$ , and $I$

First Task: Derive a number for  $v_0$  (at room temperature). We have

$$v_0 = \left( \frac{3kT}{m} \right)^{1/2} = \left( \frac{8,6 \cdot 10^{-5} \cdot 300 \text{ eV} \cdot \text{K}}{9,1 \cdot 10^{-31} \text{ K} \cdot \text{kg}} \right)^{1/2} = 1,68 \cdot 10^{14} \cdot \left( \frac{\text{eV}}{\text{kg}} \right)^{1/2}$$

- The dimension "square root of  $\text{eV/kg}$ " does not look so good - for a velocity we would like to have  $\text{m/s}$ . In looking at the energies we equated kinetic energy with the classical dimension  $[\text{kg} \cdot \text{m}^2/\text{s}^2] = [\text{J}]$  with thermal energy  $kT$  expressed in  $[\text{eV}]$ . So let's convert  $\text{eV}$  to  $\text{J}$  (use the [link](#)) and see if that solves the problem. We have  $1 \text{ eV} = 1,6 \cdot 10^{-19} \text{ J} = 1,6 \cdot 10^{-19} \text{ kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$  which gives us

$$v_0 = 1,68 \cdot 10^{14} \cdot \left( \frac{1,6 \cdot 10^{-19} \text{ kg} \cdot \text{m}^2}{\text{kg} \cdot \text{s}^2} \right)^{1/2} = 5,31 \cdot 10^4 \text{ m/s} = 1,91 \cdot 10^5 \text{ km/hr}$$

Possibly a bit surprising - those electrons are no sluggards but move around rather fast. Anyway, we have shown that a value of  $\approx 10^4 \text{ m/s}$  [as postulated in the backbone](#) is really OK.

- Of course, for  $T \rightarrow 0$ , we would have  $v_0 \rightarrow 0$  - which should worry us a bit ????. If instead of room temperature ( $T = 300 \text{ K}$ ) we would go to let's say  $1200 \text{ K}$ , we would just double the average speed of the electrons.

Second Task: Derive a number for  $\tau$ . We have

$$\tau = \frac{\sigma \cdot m}{n \cdot e^2}$$

First we need some number for the concentration of free electrons per  $\text{m}^3$ . For that we complete the [table given](#), noting that for the number of atoms per  $\text{m}^3$  we have to divide the density by the atomic weight.

Atom	Density [ $\text{kg} \cdot \text{m}^{-3}$ ]	Atomic weight $\times 1,66 \cdot 10^{-27} \text{ kg}$	Conductivity $\sigma$ $\times 10^5 [\Omega^{-1} \cdot \text{m}^{-1}]$	No. Atoms [ $\text{m}^{-3}$ ] $\times 10^{28}$
Na	970	23	2,4	2,54
Cu	8.920	64	5,9	8,40
Au	19.300	197	4,5	5,90

- So let's take  $5 \cdot 10^{28} \text{ m}^{-3}$  as a good order of magnitude guess for the number of atoms in a  $\text{m}^3$ , and for a first estimate some average value  $\sigma = 5 \cdot 10^5 [\Omega^{-1} \cdot \text{m}^{-1}]$ . We obtain

$$\tau = \frac{5 \cdot 10^5 \cdot 9,1 \cdot 10^{-31} \text{ kg} \cdot \text{m}^3}{5 \cdot 10^{28} \cdot (1,6 \cdot 10^{-19})^2 \Omega \cdot \text{m} \cdot \text{A}^2 \cdot \text{s}^2} = 3,55 \cdot 10^{-16} \frac{\text{kg} \cdot \text{m}^2}{\text{V} \cdot \text{A} \cdot \text{s}^2}$$

Well, somehow the whole thing would look much better with the unit  $[\text{s}]$ . So let's see if we can remedy the situation.

- Easy: Volts times Amperes equals *Watts* which is power, e.g. energy per time, with the unit  $[\text{J} \cdot \text{s}^{-1}] = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3}$ . Insertion yields

$$\tau = 1,42 \cdot 10^{-28} \frac{\text{kg} \cdot \text{m}^2 \cdot \text{s}^3}{\text{kg} \cdot \text{m}^2 \cdot \text{s}^2} = 3,55 \cdot 10^{-16} \text{ s} = 0,35 \text{ fs}$$

The backbone thus is right again. The scattering time is in the order of [femtosecond](#) which is a short time indeed. Since all variables enter the equation linearly, looking at somewhat other carrier densities (e.g. more than 1 electron per atom) or conductivities does not really change the general picture very much.

Third Task: Derive a number for  $v_D$ . We have (for a field strength  $E = 100 \text{ V/m} = 1 \text{ V/cm}$ )

$$\begin{aligned} |v_D| &= \frac{E \cdot e \cdot \tau}{m} = \frac{100 \cdot 1,6 \cdot 10^{-19} \cdot 3,55 \cdot 10^{-16}}{9,1 \cdot 10^{-31}} \frac{\text{V} \cdot \text{C} \cdot \text{s}}{\text{m} \cdot \text{kg}} = 6,24 \cdot 10^{-3} \frac{\text{V} \cdot \text{A} \cdot \text{s}^2}{\text{m} \cdot \text{kg}} \\ &= 6,24 \cdot 10^{-3} \frac{\text{kg} \cdot \text{m}^2 \cdot \text{s}^2}{\text{m} \cdot \text{kg} \cdot \text{s}^3} = 6,24 \cdot 10^{-3} \text{ m/s} = 6,24 \text{ mm/s} \end{aligned}$$

This is somewhat larger than the [value given in the backbone text](#).

- However - a field strength of **1 V/cm** applied to a *metal* is huge! Think about the current density  $j$  you would get if you apply **1 V** to a piece of metal **1 cm** thick.
- It is actually  $j = \sigma \cdot E = 5 \cdot 10^7 [\Omega^{-1} \cdot \text{m}^{-1}] \cdot 100 \text{ V/m} = 5 \cdot 10^9 \text{ A/m}^2 = 5 \cdot 10^5 \text{ A/cm}^2$ !
- For a more "reasonable" current density of  **$10^3 \text{ A/cm}^2$**  we have to reduce  $E$  hundredfold and then end up with  $|v_D| = \mathbf{0,0624 \text{ mm/s}}$  - and that is slow indeed!

Fourth Task: Derive a number for  $l$ . We have

$$l_{\min} = 2 \cdot v_0 \cdot \tau = 2 \cdot 5,31 \cdot 10^4 \cdot 3,55 \cdot 10^{-16} \text{ m} = 3,77 \cdot 10^7 \text{ m} = 0,0377 \text{ nm}$$

- Right again! If we add the comparatively miniscule  $v_D$ , nothing would change. Decreasing the temperature would lower  $l$  to eventually zero, or more precisely, to  $2 \cdot v_D \cdot \tau$  and thus to a value far smaller than an atom..