

## 1.15 The Legendre-Transformation in 1D

We investigate the function  $y(x)$  and  $z := dy/dx$ .

The "total differential" is

$$dy = z dx \quad . \quad (1.25)$$

Calculating

$$F(z) = y(x(z)) - zx(z) \quad (1.26)$$

we find its derivation

$$dF/dz = dy/dx(z)dx/dz(z) - x(z) - zdx/dz(z) = -x(z) \quad , \quad (1.27)$$

the "total differential" is

$$dF = -x dz \quad . \quad (1.28)$$

The transformation in equation 1.26 is called **Legendre-Transformation**. A coordinate is replaced by its force. The main advantage of this transformation is the inverse transformation (Legendre transformation of  $F(z)$ ).

We find:

$$G(x) = F(z(x)) - (-x)z(x) = y(x(z(x))) - z(x)x(z(x)) + xz(x) = y(x) \quad , \quad (1.29)$$


which is the original function **without any loss of information**.

Neglecting the additional minus sign of the inverse transformation the pair

$$\text{coordinate} \Leftrightarrow \text{force}$$

is absolutely symmetric; e.g. depending on the potential  $-p$  is a force, respectively  $p$  is a coordinate.

- Graphical representation of the Legendre transformation
- Description of the curve by the "wrapping tangents"
- for each  $x$  only one slope  $z$  must exist in order to get a well defined inverse function
- What would happen, if for a given pressure two possible volumes would exist (not possible!!, not stable!!)
- Thermodynamic functions are always strictly convex and therefore stable



- The Legendre transformation allows to transform within a potential from the intensive to the extensive parameter (and vice versa) without loss of information. This calculated potential automatically describes the corresponding thermodynamic contact correctly.
- The Legendre transformation can be applied to any coordinate independently.
- All contacts can thus be described when knowing one thermodynamic potential of the system for just one thermodynamic contact.