

3.1.2 Summary to: Polarization and Dielectric Constant

The dielectric constant ϵ_r "somehow" describes the interaction of dielectric (i.e. more or less insulating) materials and electrical fields; e.g. via the equations \Rightarrow

- \underline{D} is the **electrical displacement** or **electrical flux density**, sort of replacing \underline{E} in the Maxwell equations whenever materials are encountered.
- C is the capacity of a parallel plate capacitor (plate area A , distance d) that is "filled" with a dielectric with ϵ_r
- n is the index of refraction; a quantity that "somehow" describes how electromagnetic fields with extremely high frequency interact with matter.
in this equation it is assumed that the material has no magnetic properties at the frequency of light.

$$\underline{D} = \epsilon_0 \cdot \epsilon_r \cdot \underline{E}$$


$$C = \frac{\epsilon_0 \cdot \epsilon_r \cdot A}{d}$$

$$n = (\epsilon_r)^{1/2}$$

Electrical fields inside dielectrics polarize the material, meaning that the vector sum of electrical dipoles inside the material is no longer zero.

- The decisive quantities are the dipole moment $\underline{\mu}$, a vector, and the Polarization \underline{P} , a vector, too.
- Note: The dipole moment vector points from the negative to the positive charge - contrary to the electrical field vector!
- The dipoles to be polarized are either already present in the material (e.g. in H_2O or in ionic crystals) or are induced by the electrical field (e.g. in single atoms or covalently bonded crystals like Si)
- The dimension of the polarization \underline{P} is $[\text{C}/\text{cm}^2]$ and is indeed identical to the net charge found on unit area on the surface of a polarized dielectric.

$$\underline{\mu} = q \cdot \underline{\xi}$$

$$\underline{P} = \frac{\sum \underline{\mu}}{V}$$


The equivalent of "Ohm's law", linking current density to field strength in conductors is the Polarization law:

- The decisive material parameter is χ ("kee"), the **dielectric susceptibility**
- The "classical" flux density \underline{D} and the Polarization are linked as shown. In essence, \underline{P} only considers what happens in the material, while \underline{D} looks at the total effect: material plus the field that induces the polarization.

$$\underline{P} = \epsilon_0 \cdot \chi \cdot \underline{E}$$

$$\epsilon_r = 1 + \chi$$

$$\underline{D} = \underline{D}_0 + \underline{P} = \epsilon_0 \cdot \underline{E} + \underline{P}$$

Polarization by necessity moves masses (electrons and / or atoms) around, this will not happen arbitrarily fast.

- ϵ_r or χ thus must be functions of the frequency of the applied electrical field, and we want to consider the whole frequency range from **RF** via **HF** to light and beyond.

$\epsilon_r(\omega)$ is called the "**dielectric function**" of the material.

The tasks are:

- Identify and (quantitatively) describe the major mechanisms of polarization.
- Justify the assumed linear relationship between \underline{P} and χ .
- Derive the dielectric function for a given material.

Questionnaire

Multiple Choice questions to all of 3.1