

Radiocarbon (C14) Dating

The Basics

Advanced

Of course you know how radiocarbon or C14 dating works on principle. I did credit you with a halfway decent education, [remember](#)? So in what follows I only want to refresh your memory.

The first thing to remember is that plants turn the carbon dioxide or CO₂ contained in the air into solid organic stuff like barley (for making beer), leaves (for shading beer gardens), grass, cauliflower. Or worse, leeks, celery, Brussels sprouts and all kinds of salad. Then your wife (or kids if they study medicine like mine) force you to eat that. Fortunately animals like chicken, cows and pigs also eat the plant stuff, turning it into something useful like steaks or ham which some of us also eat.

Whatever the food chain looks like in detail, one fact applies to everything that was once alive. As soon as something organic like you dies in some (hopefully far away) time in the future, or the tree you felled with your car, or the grass that withered in the sun, there is now some dead body around that contains a lot of carbon that came exclusively from the carbon dioxide in the air via the food chain. Now something special happens. The carbon in the dead organic matter slowly changes over the years. And with carbon I mean the individual carbon *atoms*, no matter in what kind of molecules they find themselves as time goes by. When you're dead you rot and turn to compost but this involves only *chemical* reactions that change the *molecules* but not the atoms. The basic insight is that all the oxygen, hydrogen, phosphorous, nitrogen and so on atoms do not change at all - except for the carbon atoms. We need to answer two basic questions now:

1. How can an *atom* change itself? There are two basic ways:

1. The nucleus of an atom, the tiny assembly of protons and neutrons in the center of the atom, emits an elementary particle like an electron (β - decay), an helium nuclei (2 protons plus 2 neutrons; α - decay), a positron, or whatever. This is simply **radioactive decay** and what's left is a different atom.
2. The nucleus of an atom gets hit by some energetic particle like a proton or a neutron in such a way that the composition of the nucleus changes. Sort of the reverse of radioactive decay. A different atom is produced once more.

[Advanced Link](#)

Elementary particles

You might wonder about the electrons surrounding any atom - what's their role in this? None. They just rearrange to whatever is needed. The energy involved for this is minuscule compared to what's going on in the nucleus and thus can be neglected.

2. How can some carbon *atom* be different from any other carbon atom?

- It's called "*isotopes*", remember? The number of protons in a nuclei decide what element it is and thus is fixed, but the number of neutrons can be different for the same number of protons.
- The nucleus of a carbon atom, for example, has *always* 6 protons (add 6 electrons and you have a carbon atom) but the number of neutrons can be 6, 7 or 8. That gives us three carbon *isotopes*: ¹²C, ¹³C, and ¹⁴C.

[Basic Link](#)

Isotopes


The last thing to remember now is that some isotopes are stable forever, some decay radioactively. As it turns out, the carbon isotopes ¹²C and ¹³C are stable. ¹⁴C, however, decays with a half-time of 5 730 ± 40 years. In other words, if you start with 1 000 ¹⁴C atoms, you will only have 500 left after 5 730 years. 250 after 2 × 5 730 years = 11 460 years, and so on.

How It Works

All elements were made some time ago (like 10 billion years or so), either in the big bang or inside a sun. The ratio of the isotopes of a given element was fixed right then, and all the radioactive ones have long since disappeared (with a few exceptions that are of no interest here). The ratio of ¹²C to ¹³C thus is fixed at 99 : 1, and this ratio is immutable.

There is no ¹⁴C left over from the times when all the carbon atoms were made.

However, the atmosphere of our planet consists of 78.08% nitrogen molecules (N₂), 20.95% oxygen molecules (O₂), 0.93% argon (Ar), 0.038% carbon dioxide molecules (CO₂) plus traces of other stuff of no importance here. What is of importance here is that all these molecules and atoms in the upper strata of the atmosphere get hit by highly energetic particles of the cosmic radiation that bombard the earth relentlessly all the time. The atmospheric layers up there stop most of this radiation. That is generally a good thing for normal people because we would not be here or last very long if we would be hit by that radiation. It is also a good thing for archeologists because some of that radiation hits nitrogen atoms and turns them into the ¹⁴C isotope. The precise reaction is



- In words: A nitrogen atom with (per definition) 7 protons and 7 neutrons (adding up to ${}^{14}\text{N}$) gets hit by a neutron (n) that was supplied by cosmic radiation, and turns into the unstable ${}^{15}\text{N}$ isotope that decays immediately into ${}^{14}\text{C}$ by emitting a proton (p).

In yet other words: In the upper atmosphere ${}^{14}\text{C}$ is generated at a constant rate. It trickles down, forms carbon dioxide, and gets incorporated into plants and finally into everything alive. It decays inside you but as long as we live, we replenish it and essentially keep up a constant level of ${}^{14}\text{C}$ in our organic bodies. As soon as we die, no more ${}^{14}\text{C}$ is added to the remains and whatever was there now decreases in concentration by a factor of two every 5 730 years.

Radiocarbon dating now is easy. Measure the relative concentration of the ${}^{14}\text{C}$ isotope in living (or freshly dead) organic material, and you have the starting point. It is about 1 part per trillion, i.e. you have about 600 billion radioactive carbon atoms in one mol of carbon. A [mol](#) is a defined amount of stuff.

- If the organic matter you unearthed somewhere (typically coal from some old fire, bones or wood remains; all containing carbon) happens to be 5 730 years old, you would find only half of that concentration or only 300 billion radioactive carbon atoms in one mol of carbon.

With a little simple math, the "**Before Present**" or **BP** age t_{BP} of a sample comes out to

$$t_{BP} = - 5\,730 \text{ years} \cdot \log_2 \frac{N}{N_0}$$

- N is the measured ${}^{14}\text{C}$ concentration; N / N_0 thus the relative concentration of how much ${}^{14}\text{C}$ is left. This follows straight from our old [beer math](#).

Is that cool or what? So of course you know everything about the hero of science who came up with that monumental recipe for dating old organic things: **Willard Frank Libby** and his team at the University of Chicago first came up with the method in 1946; Libby got the Nobel price in 1960.

- The basics are clear now. It only remains to look at the small print coming with the general recipe for ${}^{14}\text{C}$ dating. We need to answer two big questions:

- How do you measure those extremely small amounts?
- How reliable are the results you get, or in other words: how large are the error margins and what determines their magnitude?

How It Is Done

How is it done? Well - the stuff is radioactive after all! So get a Geiger counter or whatever else registers radioactive decay, hold it over your sample, and measure how many "clicks" you get per time unit (= N). Repeat with a sample of modern carbon that contains exactly as much carbon as the sample you want to date to give you the N_0 . Use the equation form above and you are done.


- OK - that's how Libby did it, and there are problems, of course. From tiny samples you don't get that much signal, so your N value has an error of roughly $N^{1/2}$. Your N_0 value has an error - and how do you determine the exact weight you need (or just the exact carbon weight of your sample)? Then your half-time value may be a bit off. In fact, Libby used a value of 5568 ± 30 years that was later corrected to 5730 ± 40 years. Maybe your sample is contaminated with a bit of "fresh" carbon? The tree those Greek guys used to barbecue their rabbit was a 500 year old oak that was felled by a flash of lightning - you are 500 years off the proper date of the fire! An so on.

The most tricky source of error, however, is that the ${}^{14}\text{C}/{}^{12}\text{C}$ ratio for "fresh" carbon is not really constant as was assumed as a matter of course. This may have several reasons:

- Some of the cosmic radiation comes from our sun, and she is not totally reliable but wiggles the intensity a bit every now and then.
- The magnetic field of the earth influences cosmic rays somewhat (it deflects them to some degree). The magnetic field of the earth changes on large time scales and so does the ${}^{14}\text{C}/{}^{12}\text{C}$ ratio.
- The earth also exchanges carbon from the soil or from the oceans with the atmosphere at fluctuating rates, influencing the ${}^{14}\text{C}/{}^{12}\text{C}$ ratio.
- And so on. Supernova explosions nearby may increase the cosmic radiation intensity for a while or maybe

not. We just don't know.


Since we have some more methods to date things, and since more and more details become clear, we now have a calibration curve:



As the examples show, a radiocarbon age of 20 000 (10 000) years translates into 23 900 (11 400) BP years or 21 900 (9 400) BC. So be careful about the BP and the BC! With this curve we get rid of the fluctuating $^{14}\text{C}/^{12}\text{C}$ ratio problem, at least to some extent. We are also rather certain that the half-life is now as correct as it can be. The only problem that remains is to measure the amount of ^{14}C still there. That gets more difficult for old things because most of it has already decayed and thus disappeared, and for very small specimens that do not contain much to start with. The decay rates then are so low that they do not register anymore in your Geiger counter or more sophisticated equipment.

That's when "accelerator mass spectrometry" (AMS) kicks in. This is rather tricky (and very expensive) but allows a far better analysis of the $^{14}\text{C}/^{12}\text{C}$ ratio for far smaller samples than the direct method.

How does it work? You atomize your specimen, ionize the carbon (and whatever else will get ionized), accelerate those ions by high voltage, run them through a thin film to strip off electrons in order to destroy all charged molecules (like $^{13}\text{C}^1\text{H}$) that might be in there and would be confused with ^{14}C , accelerate the pure ions now once more with a really high voltage (several million Volts), and separate different atoms and their various isotopes by magnetic fields into individual ions beams. Measure how much ^{14}C and ^{12}C flies along and you have the ratio. Here is an impression of what is involved (from the Leibniz Laboratory for Radiometric Dating and Stable Isotope Research of my University (Christian-Albrechts-University of Kiel, Germany)).



Doesn't look cheap, isn't cheap, and is not all that easy to operate. You need to know exactly what you are doing, in particular with respect to specimen preparation. The data you get are only as good as your specimen. Breathe at it once in the wrong way, and it's contaminated with fresh carbon, yielding faulty results.

Not sparing time and money, how good can it be? Well, after about ten half-lives, approximately 60,000 years, not much ^{14}C is left ($1/2^{10} \approx 1/1\ 000$). That gives sort of the lower limit for a 1 mg sample. Very young stuff is problematic too, but why should you want to date that?

- Pure iron, copper, gold and so on, not containing any carbon, cannot be dated this way at all, of course. But what about steel? It contains carbon and that we can dated. Yes - but where did that carbon come from? If it came exclusively from the charcoal used in the smelting process, you get the age of the tree the charcoal was made from, and that should be close enough. However, if some of the carbon came from the limestone or calcium carbonate (CaCO_3) that was added for some reason, or if you used coal (from trees that died many million years ago) your results are useless.