

5.2 Caloric properties of ideal-gas mixtures

To calculate the chemical potential μ_i of one component i we combine Eq. (3.78) and $\mu = \Delta G_m$

$$\mu_i = \mu_i^0 + RT \ln \frac{p_i}{p^0} \quad \text{with} \quad p_i = x_i p_{tot} \quad \Rightarrow \quad \mu_i = \mu_i^0 + RT \ln x_i + RT \ln \frac{p_{tot}}{p^0} \quad (5.1)$$

This is Dalton's law. Normally p_{tot} and p^0 (the standard pressure) are selected 1 bar, thus, the second ln-term is vanishing.

For an extensive property Φ the difference by mixing compared to the pure components is

$$\Delta_{mix} \Phi^{id} = \Phi^{id}(\text{mixture}) - \sum_i n_i \Phi_m^{id}(\text{pure}) \quad (5.2)$$

Since no interaction between the ideal gas molecules exist we find

$$\Delta_{mix} H^{id} = \Delta_{mix} V^{id} = 0 \quad (5.3)$$

To calculate the effect of mixing on the entropy we apply the scheme of Fig. 5.1. Since S is a state function we can replace the calculation of mixing by expansion performed separately for each component. Since this ideal mixing is isothermal ($0 = dU = T dS - p dV$)

$$dS = \frac{p}{T} dV = \frac{n RT}{VT} dV = n R d \ln V \quad (5.4)$$

so

$$\begin{aligned} \Delta_{mix} S_1^{id} &= n_1 R \ln \frac{V_1 + V_2}{V_1}, & \Delta_{mix} S_2^{id} &= n_2 R \ln \frac{V_1 + V_2}{V_2} \\ \Rightarrow \Delta_{mix} S_1^{id} &= -n_1 R \ln x_1, & \Delta_{mix} S_2^{id} &= -n_2 R \ln x_2 \\ \Rightarrow \Delta_{mix} S^{id} &= \Delta_{mix} S_1^{id} + \Delta_{mix} S_2^{id} = - \sum_i n_i R \ln x_i \end{aligned} \quad (5.5)$$

Thus

$$\Delta_{mix} G^{id} = \sum_i n_i RT \ln x_i \quad \text{i.e.} \quad \Delta_{mix} G_m^{id} = -T \Delta_{mix} S_m^{id} = \sum_i x_i RT \ln x_i \quad (5.6)$$

Since $\Delta_{mix} S > 0$ and $\Delta_{mix} G < 0$ mixing of gases is SPONTANEOUS, i.e. an irreversible process, even for ideal gases for which no energy contributions exist.




Figure 5.1: Scheme for mixing of two ideal gases.