

5.6 Example for negative partial molar volume: salt + water

As shown in Fig. 5.3 a) partial molar volumes can also be negative, e.g. for salts which decrease the volume of a solution when added. The partial volumes represent the slope in Fig. 5.3 a). At point b, the partial molar volume is negative.

The mathematical procedure for analyzing the partial molar volume and some general explanation for the occurrence of a negative partial molar volume we will discuss using the curves of Fig. 5.4. Of course the decreasing volume (shrinking) of the mixture when adding the solute (B) is interrelated with negative values of the partial molar volumes.

- The (systematic) volume enhancement at low $b=[B]$ can be interpreted as follows: for low $[B]$ the mixture behaves almost ideal, thus there must be an increase in volume when adding B into A.
- The standard interpretation for the shrinkage is: the "open" structure of water (cf. H-bonding) is more and more disturbed, more and more water molecules are arranged with high coordination numbers around solved ions, thus leading to a volume reduction.




Figure 5.4: Volume of mixing and partial molar volume of solute B: dV/dx_b .

$V(\text{mixing})$ is experimentally treated by putting into A (solvent) variable amounts of B (solute) and measuring the volume of the mixture as shown in Fig. 5.4

1. step: The measured data $V(n_B)$ is fitted to a polynomial function

$$V_{p,T}(n_B) = a_0 + a_1 n_B + a_2 n_B^2 + a_3 n_B^3 \quad (5.14)$$

2. step: the partial molar V of B is calculated by differentiation

$$\left(\frac{\partial V}{\partial n_B}\right)_{p,T,n_A} = a_1 + 2a_2 n_B + 3a_3 n_B^2 \quad (5.15)$$

3. step: now the partial molar V of A can be calculated from

$$\begin{aligned} V &= n_A \left(\frac{\partial V}{\partial n_A}\right)_{p,T,n_B} + n_B \left(\frac{\partial V}{\partial n_B}\right)_{p,T,n_A} \\ \Rightarrow \left(\frac{\partial V}{\partial n_A}\right)_{p,T,n_A} &= \left(V - n_B \left(\frac{\partial V}{\partial n_B}\right)_{p,T,n_A}\right) / n_A \end{aligned} \quad (5.16)$$

This correlation between both partial molar volumes reflects the Gibbs-Duhem equation discussed below: the maxima are found at the same position but with opposite slopes.